

Dear Healthcare Professional,

Thank you for your unsolicited request for information. Accompanying this letter is the following information you requested on Purified Cortrophin® Gel. If we can be of any further assistance, please contact our Medical Information department at (844) CORT-GEL (844-267-8435) between the hours of 9:00 AM to 7:00 PM ET (6:00 AM to 4:00 PM PT), Monday through Friday or via email at cortrophinmedinfo@anipharmaceuticals.com.

Purified Cortrophin Gel is indicated in the following disorders:

1. Rheumatic disorders:

As adjunctive therapy for short-term administration (to tide the patient over an acute episode or exacerbation) in:

Psoriatic arthritis.

Rheumatoid arthritis, including juvenile rheumatoid arthritis (selected cases may require low-dose maintenance therapy).

Ankylosing spondylitis.

Acute gouty arthritis.

2. Collagen diseases:

During an exacerbation or as maintenance therapy in selected cases of:

Systemic lupus erythematosus.

Systemic dermatomyositis (polymyositis).

3. Dermatologic diseases:

Severe erythema multiforme (Stevens-Johnson syndrome).

Severe psoriasis.

4. Allergic states:

Atopic dermatitis

Serum sickness.

5. Ophthalmic diseases:

Severe acute and chronic allergic and inflammatory processes involving the eye and its adnexa such as:

Allergic conjunctivitis.

Keratitis.

Iritis and iridocyclitis.

Diffuse posterior uveitis and choroiditis.

Optic neuritis.

Chorioretinitis.

Anterior segment inflammation.

6. Respiratory diseases:
Symptomatic sarcoidosis.

7. Edematous states:
To induce a diuresis or a remission of proteinuria in the nephrotic syndrome without uremia of the idiopathic type or that due to lupus erythematosus.

8. Nervous system:
Acute exacerbations of multiple sclerosis.

Purified Cortrophin Gel is contraindicated for intravenous administration.

Purified Cortrophin Gel is contraindicated in patients with scleroderma, osteoporosis, systemic fungal infections, ocular herpes simplex, recent surgery, history of or the presence of a peptic ulcer, congestive heart failure, hypertension, or sensitivity to proteins derived from porcine sources.

Purified Cortrophin Gel is contraindicated in patients with primary adrenocortical insufficiency or adrenocortical hyperfunction.

Please see the enclosed Purified Cortrophin Gel Prescribing Information (PI) for detailed information including Warnings and Precautions and Adverse Reactions as well as the appropriate use of Purified Cortrophin Gel.

This communication may contain confidential, proprietary, and/or privileged information. It is intended solely for the use of the addressee. If you are not the intended recipient, you are strictly prohibited from disclosing, copying, distributing or using any of this information. If you received this communication in error, please contact the sender immediately and destroy the material in its entirety, whether electronic or hard copy.

Thank you for your inquiry.

Sincerely,



Steve Wu, PharmD
ANI Pharmaceuticals Medical Information

Potential Mechanism of Action of Purified Cortrophin[®] Gel (Repository Corticotropin Injection USP) 80U/mL in Patients With Proteinuria Due to Nephrotic Syndrome

Abstract

- This document provides summary information pertaining to Purified Cortrophin Gel (repository corticotropin injection USP) and its indication to induce a diuresis or a remission of proteinuria in nephrotic syndrome (NS) without uremia of the idiopathic type of that due to lupus erythematosus.
- It also summarizes information regarding expression of MCRs on and the potential effects on immune cells and kidney cells.

Note that this document is for information purposes only. Please refer to the Purified Cortrophin Gel (repository corticotropin injection USP) USPI for [full prescribing information](#). ANI Pharmaceuticals does not recommend the use of its products in any manner inconsistent with the FDA-approved labeling.

To report an adverse event for any ANI Pharmaceuticals product, please call 1-800-308-6755 or contact the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Email: drugsafety@anipharmaceuticals.com

Introduction

Clinical Background


Purified Cortrophin Gel™ (repository corticotropin injection USP) is approved by the FDA for use to induce a diuresis or a remission of proteinuria in nephrotic syndrome (NS) without uremia of the idiopathic type or that due to lupus erythematosus.¹

Composition of Purified Cortrophin Gel

Purified Cortrophin Gel is a porcine derived purified corticotropin, adrenocorticotrophic hormone (ACTH), in a sterile solution of gelatin. It is made up of a complex mixture of ACTH, ACTH related peptides, and other porcine pituitary derived peptides.¹

The drug product is a sterile preparation containing 80 USP units per mL and it contains 0.5% phenol (as preservative), 15.0% gelatin (for prolonged activity), water for injection, and the pH is adjusted with hydrochloric acid and sodium hydroxide.¹

Purified Cortrophin Gel contains the porcine-derived ACTH (1-39) with the following amino acid sequence¹:



Purified Cortrophin Gel Clinical Pharmacology

ACTH, the active agent in Purified Cortrophin Gel, is the anterior pituitary hormone which stimulates the functioning adrenal cortex to produce and secrete adrenocortical hormones.¹

Following administration of a single intramuscular injection of 80 units of Purified Cortrophin Gel to healthy volunteers (n=20) in an open label pharmacodynamic study, the median time (range) to reach peak plasma cortisol concentration was 8 (3 to 12) hours. The baseline corrected geometric mean maximum (CV%) cortisol levels were 34.52 µg/dL (28.2%).¹

The porcine-derived ACTH (1-39) found in Purified Cortrophin Gel is biologically similar to endogenous human ACTH,² and of the same class as other, FDA-approved natural-product and synthetic ACTH formulations.^{1,3-5}

ANI conducted a study on the pharmacodynamic effect of Purified Cortrophin Gel, including E_{max} , $AUEC_{0-24}$, and TE_{max} , and compared it with the response of the same or similar depot structures from published literature.⁶

Proposed Mechanism of Action of ACTH Potentially Related to Proteinuria Due to Nephrotic Syndrome

ANI Pharmaceuticals is not aware of any published (or unpublished) preclinical or clinical trials evaluating the mechanism of action of Purified Cortrophin Gel.

The following sections provide a brief overview of select preclinical studies and corroborative clinical observations that may help characterize the mechanism of action of ACTH and the potential role of melanocortin receptors in mediating some of its therapeutic effects.

Melanocortin Receptors and ACTH

The endogenous receptors of ACTH are the melanocortin receptors, or MCRs, of which there are 5 known isoforms (MC1R-MC5R). MCRs are broadly expressed in human tissues, including the adrenal glands, immune cells, and within glomerular substructures.^{7,8} Activation of different isoforms of MCRs may have different downstream effects depending on the tissues or cells on which they are expressed. ACTH binds to all 5 MCR isoforms, while α -MSH is a peptide derivative of ACTH that is often used to elucidate potential nonsteroidogenic actions due to its relatively low affinity for MC2R.⁹

Steroidogenic Effects of ACTH

ACTH, the active agent in Purified Cortrophin Gel, is known to stimulate production of glucocorticoids such as cortisol in the adrenal glands, a phenomenon which has been well characterized.^{10,11} This effect is attributed to ACTH agonism of MC2R, which is expressed in the adrenal cortex.¹²

MCR Expression on Immune Cells

MCRs are expressed on a number of circulating leukocytes. In human cell-based assays, these include B lymphocytes, monocytes, macrophages, granulocytes, natural killer cells, CD4⁺ T_h cells, and regulatory T cells (T_{regs}), which suggests a potential target for nonsteroidogenic stimulation by ACTH through these receptors.¹³

Potential Nonsteroidogenic Effects of ACTH on Inflammatory Cells

In cell-based assays, addition of α -MSH to lipopolysaccharide (LPS)- or TNF- α -stimulated monocyte and macrophage cultures suppressed expression of TNF- α , a proinflammatory cytokine, and activation of NF- κ B, an important pro-inflammatory mediator.^{14,15} In another cellular assay, ACTH administration reduced IgG and IgM accumulation and inhibited proliferation of activated B cells.¹⁶ In CD28 knockout mice, which are normally deficient in T_{regs}, ACTH promoted more phenotypical T_{reg} cells.¹⁷

MCR Expression on Kidney Cells

Human and animal tissue expression studies point to several MCR isoforms variously being expressed in kidney parenchyma, including podocytes, mesangial cells, tubular epithelial cells, and glomerular endothelial cells.⁷ Additionally, compared to healthy donors, the damaged glomeruli of patients with membranous nephropathy and focal segmental glomerulosclerosis have significantly elevated MC1R expression.¹⁸

Potential Nonsteroidogenic Effects of ACTH in the Kidney

In rodent models of nephrotic injury similar to that seen in NS, stimulation of MC1R-expressing podocytes through α -MSH (a nonspecific melanocortin peptide that has no appreciable affinity for MC2R) was reported to reduce proteinuria, improve glomerular morphology, and reduce oxidative stress.⁹ Other preclinical data suggest that agonism of MC1R on podocytes elicits its effects through stabilization of cytoskeletal components important to the integrity of foot processes and maintenance of the glomerular filtration barrier.^{18,19}

MCR Expression on Liver Cells

MCR gene expression has been reported in human liver tissue, most notably MC1R, suggesting activation of MCRs may have direct action on liver cells that is not dependent upon the steroidogenic pathway.²⁰

Potential Nonsteroidal Effects of ACTH in Dyslipidemia

In liver cell cultures, treatment with ACTH promotes uptake of low-density lipoprotein (LDL).²¹ Activation of MC1R by α -MSH and LD211 (a specific MC1R agonist) receptors on human hepatocytes *in vitro* has been reported to reduce cellular cholesterol content and enhance LDL and high-density lipoprotein (HDL) uptake.²²

ACTH has been shown to have a lipid-lowering effect in healthy volunteers,²³ and ACTH treatment was associated with improvement in serum lipoprotein pattern and glomerular function in patients with NS.²⁴ In patients with idiopathic membranous nephropathy, the results of a prospective trial suggested that lipid-lowering drugs may be effective in controlling proteinuria.²⁵

Citations

1. Purified Cortrophin™ Gel (Repository Corticotropin Injection USP). ANI Pharmaceuticals, Inc.; 11/21. <https://cortrophin.com/pdfs/purified-cortrophin-gel-prescribing-information.pdf>
2. Upton GV, Hollingsworth DR, Lande S, Lerner AB, Amatruda TT. Comparison of purified human and porcine ACTH in man. *The Journal of Clinical Endocrinology & Metabolism*. 1970;30(2):190-195. doi:10.1210/jcem-30-2-190
3. Atnajs Pharma UK Ltd. Synacthen Depot Ampoules 1 mg/ml. Summary of Product Characteristics (SmPC). Published October 4, 2021. Accessed November 12, 2023. <https://www.medicines.org.uk/emc/product/10823/smpc/>
4. Berkovich R, Bakshi R, Amezcua L, et al. Adrenocorticotrophic hormone versus methylprednisolone added to interferon β in patients with multiple sclerosis experiencing breakthrough disease: a randomized, rater-blinded trial. *Ther Adv Neurol Disord*. 2017;10(1):3-17. doi:10.1177/1756285616670060

5. *ACTHAR GEL (Repository Corticotropin Injection), for Intramuscular or Subcutaneous Use. Prescribing Information.* Mallinkrodt ARD LLC; 2021. Accessed November 12, 2023. <https://acthar.com/Static/pdf/Acthar-PI.pdf>
6. ANI Pharmaceuticals, Inc. Data on file.
7. Gong R. Leveraging melanocortin pathways to treat glomerular diseases. *Adv Chronic Kidney Dis.* 2014;21(2):134-151. doi:10.1053/j.ackd.2013.09.004
8. Chang M, Chen B, Shaffner J, Dworkin LD, Gong R. Melanocortin system in kidney homeostasis and disease: novel therapeutic opportunities. *Front Physiol.* 2021;12:651236. doi:10.3389/fphys.2021.651236
9. Lindskog A, Ebefors K, Johansson ME, et al. Melanocortin 1 receptor agonists reduce proteinuria. *J Am Soc Nephrol.* 2010;21(8):1290-1298. doi:10.1681/ASN.2009101025
10. Nussey S, Whitehead S. *Endocrinology: An Integrated Approach.* BIOS Scientific Publishers; 2001. Accessed January 26, 2022. <http://www.ncbi.nlm.nih.gov/books/NBK22/>
11. Jenkins D, Forsham PH, Laidlaw JC, Reddy WJ, Thorn GW. Use of ACTH in the diagnosis of adrenal cortical insufficiency. *Am J Med.* 1955;18(1):3-14. doi:10.1016/0002-9343(55)90200-x
12. Novoselova TV, King PJ, Guasti L, Metherell LA, Clark AJL, Chan LF. ACTH signalling and adrenal development: lessons from mouse models. *Endocr Connect.* 2019;8(7):R122-R130. doi:10.1530/EC-19-0190
13. Gong R. The renaissance of corticotropin therapy in proteinuric nephropathies. *Nat Rev Nephrol.* 2011;8(2):122-128. doi:10.1038/nrneph.2011.190
14. Yoon SW, Goh SH, Chun JS, et al. α -Melanocyte-stimulating hormone inhibits lipopolysaccharide-induced tumor necrosis factor- α production in leukocytes by modulating protein kinase A, p38 kinase, and nuclear factor κ B signaling pathways. *Journal of Biological Chemistry.* 2003;278(35):32914-32920. doi:10.1074/jbc.M302444200
15. Manna SK, Aggarwal BB. α -melanocyte-stimulating hormone inhibits the nuclear transcription factor NF- κ B activation induced by various inflammatory agents. *The Journal of Immunology.* 1998;161(6):2873-2880. doi:10.4049/jimmunol.161.6.2873
16. Olsen NJ, Decker DA, Higgins P, et al. Direct effects of HP Acthar Gel[®] on human B lymphocyte activation in vitro. *Arthritis Res Ther.* 2015;17:300. doi:10.1186/s13075-015-0823-y
17. Zhao J, Jiang L, Uehara M, et al. ACTH treatment promotes murine cardiac allograft acceptance. *JCI Insight.* 2021;6(13):e143385. doi:10.1172/jci.insight.143385
18. Bergwall L, Wallentin H, Elvin J, et al. Amplification of the melanocortin-1 receptor in nephrotic syndrome identifies a target for podocyte cytoskeleton stabilization. *Sci Rep.* 2018;8(1):15731. doi:10.1038/s41598-018-34004-7

19. Elvin J, Buvall L, Lindskog Jonsson A, et al. Melanocortin 1 receptor agonist protects podocytes through catalase and RhoA activation. *Am J Physiol Renal Physiol*. 2016;310(9):F846-856. doi:10.1152/ajprenal.00231.2015
20. Gatti S, Colombo G, Turcatti F, et al. Reduced expression of the melanocortin-1 receptor in human liver during brain death. *Neuroimmunomodulation*. 2006;13(1):51-55. doi:10.1159/000094513
21. Bartens W, Krämer-Guth A, Wanner C. Corticotropin increases the receptor-specific uptake of native low-density lipoprotein (LDL)—but not of oxidized LDL and native or oxidized lipoprotein(a) [LP(a)]—in HEPG2 cells: No evidence for Lp(a) catabolism via the LDL-receptor. *Metabolism*. 1997;46(7):726-729. doi:10.1016/S0026-0495(97)90113-X
22. Thapa K, Kadiri JJ, Saukkonen K, et al. Melanocortin 1 receptor regulates cholesterol and bile acid metabolism in the liver. Quinn MA, Isales C, Sánchez-Ceinos J, eds. *eLife*. 2023;12:e84782. doi:10.7554/eLife.84782
23. Arnadottir M, Berg AL, Kronenberg F, et al. Corticotropin-induced reduction of plasma lipoprotein(a) concentrations in healthy individuals and hemodialysis patients: relation to apolipoprotein(a) size polymorphism. *Metabolism*. 1999;48(3):342-346. doi:10.1016/s0026-0495(99)90083-5
24. Berg AL, Nilsson-Ehle P, Arnadottir M. Beneficial effects of ACTH on the serum lipoprotein profile and glomerular function in patients with membranous nephropathy. *Kidney Int*. 1999;56(4):1534-1543. doi:10.1046/j.1523-1755.1999.00675.x
25. Rayner BL, Byrne MJ, van Zyl Smit R. A prospective clinical trial comparing the treatment of idiopathic membranous nephropathy and nephrotic syndrome with simvastatin and diet, versus diet alone. *Clin Nephrol*. 1996;46(4):219-224.